



Heads Up to Schools: KNOW YOUR CONCUSSION ABCs

Assess
the
situation

Be alert for
signs and
symptoms

Contact a
health care
professional



Signs and Symptoms of a Concussion

A concussion is caused by a bump, blow, or jolt to the head. Concussions can also occur from a fall or blow to the body that causes the head to move rapidly back and forth. Even what seems to be a mild bump to the head can be serious. Be alert for **any** of the following signs and symptoms.

SIGNS OBSERVED BY SCHOOL PROFESSIONALS

- Appears dazed or stunned
- Is confused about events
- Answers questions slowly
- Repeats questions
- Can't recall events *prior* to hit, bump, or fall
- Can't recall events *after* hit, bump, or fall
- Loses consciousness (even briefly)
- Shows behavior or personality changes
- Forgets class schedule or assignments

SYMPTOMS REPORTED BY THE STUDENT

Thinking/Remembering

- Difficulty thinking clearly
- Difficulty concentrating or remembering
- Feeling more slowed down
- Feeling sluggish, hazy, foggy, or groggy

Physical

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Fatigue or feeling tired
- Blurry or double vision
- Sensitivity to light or noise
- Numbness or tingling
- Does not "feel right"

Emotional

- Irritable
- Sad
- More emotional than usual
- Nervous

Sleep*

- Drowsy
- Sleeps *less* than usual
- Sleeps *more* than usual
- Has trouble falling asleep

**Only ask about sleep symptoms if the injury occurred on a prior day.*

What can school professionals do?



Know your Concussion ABCs:

A—Assess the situation

B—Be alert for signs and symptoms

C—Contact a health care professional

**For more information contact us:*

HeadsUP! is a programme of the ComaCARE Trust
Registered Charity NPO 061-592 / Brain Injury Support Group
Janine Filmer: 021 531 0678 / 084 514 2404
Bernadette Medder: 021 794 4150 / 082 412 3333
Email: headsupsouth@gmail.com

THANKS TO U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION

